

Yaroslavl State University
 Preprint YARU-HE-97/09
 hep-ph/9712423

Field-induced axion decay $a \rightarrow e^+e^-$ in KSVZ-model

N.V. Mikheev, O.S. Ovchinnikov

Yaroslavl P.G. Demidov State University, Yaroslavl 150000, Russia
E-mail: mikheev@yars.free.net owoles@univ.uniyar.ac.ru

and

L.A. Vassilevskaya

Moscow M.V. Lomonosov State University, V-952, Moscow 117234, Russia
E-mail: vasilevs@vitep5.itep.ru

Abstract

The axion decay into electron-positron pair $a \rightarrow e^+e^-$ is studied in an external magnetic field in KSVZ-model where axions have only induced coupling to leptons. The axion lifetime in the field decreases with energy and field strength to seconds.

*Talk given at the International Workshop
 on Particle Physics and the Early Universe "COSMO'97",
 Ambleside, Lake District, England, 15-19 September 1997.*

Peccei-Quinn (PQ) symmetry $U_{PQ}(1)$ [1], with its accompanying axion [2], continues to be an attractive solution to the strong CP problem in QCD. At present axions are of great interest not only in theoretical aspects of elementary particle physics, but in some astrophysical and cosmological applications as well [3, 4, 5]. Although the original axion is excluded experimentally, modified PQ models with very light and very weakly coupled, so called, invisible axions are still tenable (see, for example the recent review [6]). Invisible axion models are classified into two types depending on whether or not they have direct couplings to leptons. The, so called, KSVZ axions [7] are hadronic axions with only induced coupling to leptons. The, so called, DFSZ axions [8], arise in models where axions couple to leptons already at tree level.

In this paper we analyze the influence of an external magnetic field on the axion decay into electron-positron pair via a photon intermediate state $a \rightarrow \gamma \rightarrow e^+e^-$ in KSVZ model. The lagrangian describing the effective axion-photon coupling is:

$$\mathcal{L}_{a\gamma} = g_{a\gamma} \partial_\mu A_\nu \tilde{F}_{\nu\mu} a, \quad (1)$$

$g_{a\gamma}$ is a constant with the dimension $(energy)^{-1}$; A_μ is the 4-potential of the quantized electromagnetic field, $\tilde{F}$ is the dual external field tensor. In the second order of the perturbation theory a matrix element can be presented in the form:

$$\begin{aligned} S &= -\frac{ie g_{a\gamma}}{\sqrt{2E_a V}} \int d^4x \left(\bar{\psi}(x) \hat{h} \psi(x) \right) e^{-iqx}, \\ h_\alpha &= (q\tilde{F}G(q))_\alpha = q_\mu \tilde{F}_{\mu\nu} G(q)_{\nu\alpha}, \end{aligned} \quad (2)$$

where $e > 0$ is the elementary charge; $\psi(x)$ is the known solution of the Dirac equation in a magnetic field; q_α is the 4-momentum of the decaying axion; $G_{\alpha\beta}$ is the photon propagator in the field. It is convenient to use the expression for the propagator in a diagonal form [9]:

$$G_{\alpha\beta} = \sum_{\lambda=1}^3 \frac{b_\alpha^{(\lambda)} b_\beta^{(\lambda)}}{(b^{(\lambda)})^2} \frac{-i}{q^2 - \mathfrak{x}^{(\lambda)}}, \quad b_\alpha^{(\lambda)} b_\alpha^{(\lambda')} = \delta_{\lambda\lambda'} (b_\alpha^{(\lambda)})^2. \quad (3)$$

Here the basis vectors $b_\alpha^{(\lambda)}$ are the eigenvectors of the photon polarization tensor, $\mathfrak{a}^{(\lambda)}$ are the eigenvalues. Notice that only the basis vector $b_\alpha^{(2)} = (q\tilde{F})_\alpha$ gives a contribution to the decay.

Here we present the results of our calculations for the axion lifetime in the limiting case $E_a^2 \sin^2 \theta \gg eB$ (E_a is the energy of the decaying axion; θ is the angle between the vectors of the magnetic field strength $\vec{B}$ and the momentum of the axion $\vec{q}$), when electron and positron are born in the states corresponding to the highest Landau levels. In this case the eigenvalue $\mathfrak{a}^{(2)}$ is described by the expression:

$$\mathfrak{a}^{(2)} \simeq \frac{9 \cdot 3^{1/6} \Gamma^4(\frac{2}{3})}{14\pi^2} \alpha (1 - i\sqrt{3}) (e^2 q F F q)^{1/3}. \quad (4)$$

To obtain the decay probability one has to carry out a non-trivial integration over the phase space of the electron-positron pair taking their specific kinematics in the magnetic field into account. The result can be presented in the form:

$$W \simeq 2,46 \cdot 10^3 g_{a\gamma}^2 (eB)^{4/3} E_a^{1/3} \sin^{4/3} \theta.$$

The corresponding expression for the axion lifetime is:

$$\tau^{KSVZ} \simeq 1,16 \left(\frac{10^{-10}}{g_{a\gamma} \text{ GeV}} \right)^2 \left(\frac{E_a}{10 \text{ MeV}} \right)^{-1/3} \left(\frac{10^{15} G}{B \sin \theta} \right)^{4/3} s. \quad (5)$$

For comparison let us present here the expression for the axion lifetime in vacuum [4]:

$$\tau^{(0)}(a \rightarrow 2\gamma) \sim 1,3 \cdot 10^{44} \left(\frac{10^{-10}}{g_{a\gamma} \text{ GeV}} \right)^2 \left(\frac{10^{-3} \text{ eV}}{m_a} \right)^4 \left(\frac{E_a}{10 \text{ MeV}} \right) s. \quad (6)$$

The work was supported by Grant INTAS 96-0659.

References

- [1] R.D. Peccei and H.R. Quinn, *Phys. Rev. Lett.* **38**, 1440 (1977); *Phys. Rev. D* **16**, 1791 (1977).
- [2] S. Weinberg, *Phys. Rev. Lett.* **40**, 223 (1978);
F. Wilczek, *Phys. Rev. Lett.* **40**, 271 (1978).
- [3] M.S. Turner, *Phys. Rept.* **197**, 67 (1990).
- [4] G.G. Raffelt, *Phys. Rept.* **198**, 1 (1990).
- [5] G.G. Raffelt, *Stars as Laboratories for Fundamental Physics* (University of Chicago Press: Chicago, 1996).
- [6] R.D. Peccei, Preprint hep-ph/9606475 (QCD, Strong CP and Axions).
- [7] J. Kim, *Phys. Rev. Lett.* **43**, 103 (1979);
M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, *Nucl. Phys. B* **166**, 493 (1980).
- [8] M. Dine, W. Fischler and M. Srednicki, *Phys. Lett. B* **104**, 199 (1981);
A.P. Zhitnisky, *Sov. J. Nucl. Phys.* **31**, 260 (1980).
- [9] A.E. Shabad, *Ann. Phys. N.Y.* **90**, 166 (1975).